

CELL ELECTROPHYSIOLOGY

**BRAIN SLICE ELECTROPHYSIOLOGY**

IN VIVO BRAIN ELECTROPHYSIOLOGY

IN VIVO SC & DRG ELECTROPHYSIOLOGY

CALCIUM IMAGING


# Calcium imaging in primary rat neuron cultures



# GLUTAMATE ELICITS CALCIUM FLUXES IN RAT PYRAMIDAL NEURONS

## METHODS


- Neuron cultures were prepared from E18 rat cortex and plated on a monolayer of astrocytes growing on PDL-coated glass imaging dishes (MatTek).
- Calcium imaging was performed in Fura-2 AM loaded cells at 9 days *in vitro*.




*Perfusion with 100  $\mu$ M glutamate elicited an influx of calcium in E18 pyramidal neurons.*

# GLUTAMATE ELICITS CALCIUM FLUXES IN RAT PYRAMIDAL NEURONS

**Baseline**




**100  $\mu$ M glutamate**



# GLUTAMATE ELICITS CALCIUM FLUXES IN RAT PYRAMIDAL NEURONS

## METHODS

- Neuron cultures were prepared from E18 rat cortex and plated on a monolayer of astrocytes growing on PDL-coated glass imaging dishes (MatTek).
- Calcium imaging was performed in Fura-2 AM loaded cells at 9 days *in vitro*.




*Calcium responses to glutamate in E18 pyramidal neurons were concentration dependent. Depolarization by 60 mM KCl also elicited a comparable calcium signal.*



# CAPSAICIN ELICITS CALCIUM RESPONSES IN RAT DRG SENSORY NEURONS

## METHODS

- Sensory neuron cultures were prepared from adult rat dorsal root ganglia and plated on a monolayer of astrocytes growing on PDL-coated glass imaging dishes (MatTek).
- Calcium imaging was performed on Fura-2 AM loaded cells at 9 days *in vitro*.




*Phase-contrast (left) and ratiometric images of Fura-2 loaded DRG sensory neurons. 300 nM capsaicin (TrpV1 agonist) elicited calcium responses in ~50% of sensory neurons.*

# CAPSAICIN ELICITS CALCIUM RESPONSES IN RAT DRG SENSORY NEURONS

## METHODS

- Sensory neuron cultures were prepared from adult rat dorsal root ganglia and plated on a monolayer of astrocytes growing on PDL-coated glass imaging dishes (MatTek).
- Calcium imaging was performed on Fura-2 AM loaded cells at 9 days *in vitro*.



*Time course of calcium responses in individual Fura-2AM loaded DRG sensory neurons indicated by colored lines.*

*Following a 2 min baseline period, perfusion for 1 min with 300 nM capsaicin (TrpV1 agonist) elicited calcium responses in ~50% of neurons expressing TrpV1. All cells responded to depolarization by 60 mM KCl indicating healthy, electrically excitable neurons.*




# Experiment #9 (12/6) DRG #8 (2 div)

## Capsaicin conc-response

### Results


- Capsaicin elicited calcium responses in rat DRG sensory neurons in a concentration-dependent fashion
- The percent of responsive cells increased with increasing concentrations of capsaicin
- The magnitude calcium response ( $\Delta R/R$ ) increased with increasing concentrations of capsaicin
- Maximum responses were seen  $\geq 300$  nM



	30 nM	100 nM	300 nM	1 uM
% responders ( $\Delta R/R > 50\%$ )	1% (1/84)	35% (42/119)	55% (46/84)	82% (64/78)
$\Delta R/R$	24%	67%	148%	130%

# AITC elicits calcium response in rat DRG sensory neurons

- 50  $\mu$ M AITC (1 min) elicits a calcium response in  $\geq 50\%$  of DRG sensory neurons
- 60 mM KCl used as positive control to identify healthy, excitable DRG sensory neurons





# Contact us



**Raymond Price, PhD, EMBA**  
**Chief Business Officer**

[Raymond.price@neuroservices-alliance.com](mailto:Raymond.price@neuroservices-alliance.com)

Mobile - +1 (858) 649 9403



**Bob Petroski, PhD**  
**CSO, Cell Electrophysiology**

[Bob.petroski@neuroservices-alliance.com](mailto:Bob.petroski@neuroservices-alliance.com)

Mobile - +1 (858) 774 4485



# The CNS Electrophysiology CRO

[WWW.NEUROSERVICES-ALLIANCE.COM](http://WWW.NEUROSERVICES-ALLIANCE.COM)